



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

# A DIRECT DEFINITION OF LOGARITHMIC DERIVATIVE.\*

By E. R. HEDRICK, University of Missouri.

**1. The Usual Definitions.** The logarithmic derivative of a function  $y = f(x)$  is usually defined either as the derivative of the Napierian logarithm of  $y$ :

$$(1) \quad \frac{d \log_e y}{dx} = \frac{d \log_e f(x)}{dx};$$

or by means of the resulting formula:

$$(2) \quad \frac{dy}{dx} \div y = f'(x) \div f(x).$$

These definitions suffer from the fact that the logarithm is defined only when  $f(x)$  is positive, though (2) may be used independently of (1) when  $f(x)$  is negative.

**2. A Direct Definition.** The importance of logarithmic differentiation in elementary work arises from its meaning as the *relative rate of increase* of  $y$  with respect to  $x$ . From this standpoint, the intervention of logarithms in the definition is wholly accidental and extraneous.

In order to define directly the relative rate of increase, we may divide the actual increase  $\Delta y$  by the increase in  $x$ ,  $\Delta x$ , divide this ratio  $\Delta y/\Delta x$  by some average value of  $f(x)$  in the range  $\Delta x$ , and take the limit of this quotient as  $\Delta x$  approaches zero. Denoting the relative rate of increase by  $r_r$ , we may write

$$(3) \quad r_r = \lim_{\Delta x=0} \left\{ \frac{\Delta y}{\Delta x} \div [\text{Average Value of } f(x)] \right\}.$$

The average value of  $f(x)$  to be used may be selected in a variety of ways; that which seems most appropriate is the usual expression:

$$[\text{Average Value of } f(x)] = \frac{1}{\Delta x} \int_{x=a}^{x=a+\Delta x} f(x)dx,$$

where  $x = a$  is the value of  $x$  at which the relative rate is to be found. Then (3) becomes

$$r_r = \lim_{\Delta x=0} \left\{ \frac{\Delta y}{\Delta x} \div \left[ \frac{1}{\Delta x} \int_{x=a}^{x=a+\Delta x} f(x)dx \right] \right\},$$

whence, simplifying and replacing  $\Delta y$  by  $f(a + \Delta x) - f(a)$ , we obtain

$$(4) \quad r_r = \lim_{\Delta x=0} \left[ \frac{f(a + \Delta x) - f(a)}{\int_a^{a+\Delta x} f(x)dx} \right].$$

---

\* Read before the Southwestern Section of the American Mathematical Society, November, 1912.

Let  $\phi(x)$  be any indefinite integral of  $f(x)$ ; then

$$\int f(x)dx = \phi(x), \quad \int_a^{a+\Delta x} f(x)dx = \phi(a + \Delta x) - \phi(a),$$

and (4) becomes

$$(5) \quad r_r = \lim_{\Delta x \rightarrow 0} \left[ \frac{f(a + \Delta x) - f(a)}{\phi(a + \Delta x) - \phi(a)} \right],$$

which we shall adopt as *the definition of the relative rate of increase of  $f(x)$ , or the logarithmic derivative of  $f(x)$ , with respect to  $x$ .*

**3. Identification with the Usual Definition.** At least when  $f(x)$  is positive, the definition (5) coincides with the usual definition (1); and (5) coincides with the definition (2) whenever (2) has a meaning.

For, since

$$(6) \quad \frac{f(a + \Delta x) - f(a)}{\Delta x} = \frac{f(a + \Delta x) - f(a)}{\phi(a + \Delta x) - \phi(a)} \cdot \frac{\phi(a + \Delta x) - \phi(a)}{\Delta x}$$

and since the limit of  $[\phi(a + \Delta x) - \phi(a)]/\Delta x$  surely exists and is equal to  $f(a)$ , if  $f(x)$  is continuous at  $x = a$ , we have

$$(7) \quad f'(a) = r_r \cdot f(a)$$

provided  $f'(a)$  exists.

Conversely, if  $r_r$ , as defined by (5), exists, and if  $f(x)$  is continuous at  $x = a$ , it follows from (6) that  $f'(a)$  exists.

**4. The Law of the Mean.** If  $f(x)$  is continuous, and if  $r_r$  exists, the derivative  $f'(x)$  exists; moreover  $\phi'(x)$  exists and is equal to  $f(x)$ ; hence, by the generalized law of the mean,\*

$$(8) \quad \frac{f(a + \Delta x) - f(a)}{\phi(a + \Delta x) - \phi(a)} = \frac{f'(c)}{\phi'(c)} = \frac{f'(c)}{f(c)},$$

where  $c$  lies between  $a$  and  $a + \Delta x$ .

The expression on the left,  $[f(a + \Delta x) - f(a)] \div [\phi(a + \Delta x) - \phi(a)]$ , may be called the average relative rate over the interval  $\Delta x$ ; with this notation, the equation (8) may be stated as follows: *The average relative rate over the interval  $\Delta x$  is precisely equal to the instantaneous relative rate at some point in that interval.*

This statement, which is the analogon to the law of the mean for ordinary derivatives, may be used to prove directly several important theorems.

Thus, if the logarithmic derivative is zero identically, it follows from (8) that  $f(x)$  is a constant.

Again, suppose that the logarithmic derivatives of two functions  $f(x)$  and  $F(x)$  are identically equal; then

$$0 = \frac{f'(x)}{f(x)} - \frac{F'(x)}{F(x)} = \frac{f'(x)F(x) - F'(x)f(x)}{[F(x)]^2} \div \frac{f(x)}{F(x)}.$$

---

\* See, for example, Goursat-Hedrick, *Mathematical Analysis*, Vol. I, p. 8; it is vital that  $c$  is the same in numerator and denominator.

The extreme right-hand member is, however, the logarithmic derivative of  $f(x)/F(x)$ ; hence by the preceding theorem,  $f(x)/F(x)$  is constant, or  $f(x) = k \cdot F(x)$ : *If two functions have the same logarithmic derivative, their quotient is a constant; or If the logarithmic derivative of a function is given, that function is determined except for a constant factor.*

These are the fundamental theorems concerning logarithmic derivatives; they are precisely analogous to the corresponding theorems for ordinary derivatives.

**5. Discontinuity.** It has been shown above that  $f'(x)$  exists whenever the relative rate  $r_r$  defined by (5) exists, provided only that  $f(x)$  is continuous. It is, however, possible that  $f'(x)$  and therefore also  $r_r$  is discontinuous.

It might appear that (8) precludes such a possibility, for the left side approaches  $r_r$ , and  $c$  approaches  $a$  as  $\Delta x$  approaches zero. But  $c$  does not necessarily take on all values; hence the approach of  $c$  to  $a$  may be only through a set of special values of  $x$ .

That the possibility just mentioned actually does occur is manifested by the example

$$f(x) = 4x^3 \sin \frac{1}{x} - x^2 \cos \frac{1}{x} + 1, \quad x \neq 0; \quad f(0) = 1;$$

which has an indefinite integral:

$$\phi(x) = x^4 \sin \frac{1}{x} + x, \quad x \neq 0; \quad \phi(0) = 0.$$

Its derivative is

$$f'(x) = 12x^2 \sin \frac{1}{x} - 5x \cos \frac{1}{x} + \sin \frac{1}{x}, \quad x \neq 0; \quad f'(0) = 0;$$

which is discontinuous. Its logarithmic derivative may be derived either by (2) or by (5), and its value when  $x = 0$  is zero.

**6. Conclusion.** The facts and illustrations given above parallel completely the fundamental theorems and illustrations usually given for ordinary derivatives. In deducing them, although the name *logarithmic* derivative has been used, the notion of logarithms has not been employed, and it has been shown that the ideas themselves are independent of the concept of logarithms.

## TWO GEOMETRICAL APPLICATIONS OF THE METHOD OF LEAST SQUARES.

By J. L. COOLIDGE, Harvard University.

The two problems discussed in the present paper are taken from a recent work by Vahlen entitled "Konstruktionen und Approximationen."\* It seems to me that they are sufficiently interesting and important to merit special attention.

\* Leipzig, 1911, pages 125, 126.